arXiv:astro-ph/0207280v1 12 Jul 2002

KILOHERTZ QUASI-PERIODIC OSCILLATIONS AND STRONG FIELD GRAVITY IN X-RAY BINARIES

MARIANO MÉNDEZ

Astronomical Institute, University of Amsterdam, the Netherlands and SRON, National Institute for Space Research, Utrecht, the Netherlands. E-mail: M.Mendez@SRON.nl

In the past five years observations with the *Rossi X-ray Timing Explorer* have revealed fast quasi-periodic oscillations in the X-ray flux of about 20 X-ray binaries. Thought to originate close to the surface of a neutron star, these oscillations provide unique information about the strong gravitational field in which they are produced.

The dynamical timescale close to an object of mass M and radius R is $\tau \sim R/v_{\rm ff} = \sqrt{R^3/2GM}$, where $v_{\rm ff}$ is the local free-fall velocity. For typical masses and radii of neutron stars ($M \sim 1-2\,{\rm M}_\odot$ and $R \sim 10-20\,{\rm km}$), $\tau \lesssim 10^{-3}$ s. It was only after NASA's *Rossi X-ray Timing Explorer (RXTE)* was launched in December 1995 that we could finally probe these short timescales, both in neutron-star and black-hole X-ray binaries. Groups at Goddard Space Flight Center and the University of Amsterdam, almost simultaneously announced the detection of submillisecond quasi-periodic variability in the X-ray flux of two binary systems with neutron star primary[1,2]. Since then, so-called kilohertz quasi-periodic oscillations (kHz QPOs) have been observed in more than 20 X-ray binary systems[3].

KHz QPOs usually appear in pairs[4], with frequencies ν_1 and $\nu_2 (> \nu_1)$ between ~ 300 Hz[5] (the lowest value for ν_1) and ~ 1330 Hz[6] (the highest value for ν_2). For a given source both ν_1 and ν_2 can drift by ~ 100 Hz on timescales of a few hundred seconds[7], with a typical dynamic range of $\sim 200-300$ Hz; up to ~ 800 Hz variations have been observed in one source[6]. Despite these rather large ν_1, ν_2 shifts, for each source $\Delta\nu = \nu_2 - \nu_1$, remains more or less constant[4,7,8]. In the best studied sources, $\Delta\nu$ has been observed to decrease by $50-100$ Hz as ν_1, ν_2 increase[9,10]. In all the other sources of kHz QPO, $\Delta\nu$ is consistent with this trend[11]

It is widely accepted that at least one of the two QPO peaks is produced by matter in Keplerian orbits around the neutron star at the inner edge of the accretion disc[12−14]. If this is so, QPO frequencies should depend sensitively upon the mass and angular momentum of the compact star as well as upon the orbital radius at which the QPOs are produced[12,13]. In principle it is possible to measure some of these parameters independently[23−25], which would allow us to set simultaneous constraints to the mass and radius of the star, and would provide means of knowing the high-density equation of state of neutronic matter[12], an issue of central importance in nuclear physics research.

On timescales of a few hours QPO frequencies correlate with X-ray intensity[4], but this correlation breaks down on timescales of a day or more[18]; on longer timescales QPO frequencies correlate much better with the X-ray spectral properties of the source[6,10−16]. It is thought that mass accretion rate is the mechanism behind this behavior[12].

The Fourier power spectra of X-ray binaries often show a broad-band component that is roughly flat below a break frequency at $\nu_{\rm b} \sim 1-10$ Hz and decreases above

ν_b, and a low-frequency QPO at $\sim 10-50$ Hz. Studies with RXTE have recently shown that both ν_b and the frequency of the low-frequency QPO are strongly correlated to the frequency of the kHz QPOs[16,26–28] and to each other[29], and that these correlation encompasses systems both with neutron star and black hole primaries. This result is quite significant, as it seems to indicate that the presence of a solid surface, a magnetic field, or an event horizon plays no significant role in the mechanism that produces the rapid variability observed in these objects.

Acknowledgments

This work was supported by the Netherlands Organization for Scientific Research, grant PGS 78-277, the Netherlands Foundation for research in astronomy, grant 781-76-017, the Netherlands Research School for Astronomy, and by LKBF. The author is grateful to Max-Planck-Institut für Astrophysik for their hospitality.

References

1. M. van der Klis et al., IAUC **6319**, 1 (1996)
2. T. Strohmayer et al., IAUC **6320**, 1 (1996)
3. M. van der Klis, ARA&A **38**, 717 (2000)
4. E. C. Ford et al., ApJ **475**, L123 (1997)
5. P. G. Jonker et al., ApJ **537**, 374 (2000)
6. S. van Straaten et al., ApJ **540**, 1049 (2000)
7. T. Strohmayer et al., ApJ **469**, L9 (1996)
8. R. Wijnands, ApJ **490**, L157 (1998)
9. M. van der Klis et al., ApJ **481**, L97 (1997)
10. M. Méndez et al., ApJ **505**, L23 (1998)
11. D. Psaltis et al., ApJ **501**, 95 (1998)
12. M. C. Miller, ApJ **508**, 791 (1998)
13. L. Stella & M. Vietri, PRL **82**, 17 (1999)
14. V. Osherovich & L. Titarchuk, ApJ , (1999)
15. T. Strohmayer et al. in *The Active X-ray Sky*, eds. L. Scarsi et al., 129 (1998)
16. E. C. Ford, ApJ **519**, L73 (1999)
17. N. Weinberg et al., ApJ , in press (2000)
18. W. Zhang et al., ApJ **495**, L9 (1998)
19. E.C. Ford et al., ApJ **486**, L47 (1997)
20. P. Kaaret et al., ApJ **497**, L93 (1998)
21. M. Méndez et al., ApJ **511**, L49 (1999)
22. M. Méndez & M. van der Klis, ApJ **517**, L51 (1999)
23. P. Kaaret et al., ApJ **520**, L37 (1999)
24. M. Méndez in *19th Texas Symp.*, eds. J. Paul, et al., 15/16 (2000)
25. P. Reig, ApJ **530**, L16 (2000)
26. T. Di Salvo, ApJ , in press (2000)
27. E. C. Ford & M. van der Klis, ApJ **506**, L39 (1998)
28. D. Psaltis et al., ApJ **520**, 262 (1999)
29. R. Wijnands & M. van der Klis, ApJ **514**, 939 (1999)